



DEPARTMENT OF HEALTH
DEPARTEMENT VAN GESONDHEID

UMNYANGO WEZEMPILO
LEFAPHA LA MAPHELO

Facsimile: 012 395 8824
Telephone: 012 395 9638
e-mail: SteelG@health.gov.za

Private Bag X828
Pretoria 0001

Enquiries: Mr G Steel
Reference: NEML15122015

NOTICE OF REQUEST FOR COMMENT ON THE STANDARD TREATMENT GUIDELINES AND ESSENTIAL MEDICINES LIST FOR PRIMARY HEALTH LEVEL OF CARE (2014 EDITION)

The ministerially appointed National Essential Medicines List (EML) Committee will start with the review of the Standard Treatment Guidelines and Essential Medicines List for Primary Health level of care, 2014 edition.

The Primary Health Level Standard Treatment Guidelines and Essential Medicines List are aimed for use by doctors and nurse prescribers providing care at primary healthcare facilities to provide access to pharmaceuticals to manage common conditions at this level.

Kindly circulate the request to relevant healthcare professionals at your institutions for comment. Constructive comment with regard to the identification of gross errors, particularly diagnosis and treatment, will be appreciated. A short motivation to be included to substantiate any comment made.

Where an alternative medicine is recommended, this should be supported by appropriate evidence. Attached is the guideline for the Motivation of a New Medicine on the National Essential Medicines List.

It would be appreciated if comments can be received by **28 February 2016**.

Comments may be submitted via fax, e-mail or by post to:

Trudy Leong
Tel: 012 395 8287
Fax to e-mail: 0862484875
E-mail: leongt@health.gov.za

Essential Drugs Programme
Private Bag X828
PRETORIA
0001

Your co-operation in this regard is appreciated.

Kind regards

pp **DIRECTOR-GENERAL**
DATE: 15 December 2015

GUIDELINES FOR THE MOTIVATION OF A NEW MEDICINE ON THE NATIONAL ESSENTIAL MEDICINES LIST

Section 1: Medication details

- » Generic name
A fundamental principle of the Essential Drug Programme is that of generic prescribing. Most clinical trials are conducted using the generic name.
- » Proposed indication
There will usually be many registered indications for the medication. However, this section should be limited to the main indication which is supported by the evidence provided in section 2.
- » Prevalence of the condition in South Africa
This information is not always readily available. However, it is an important consideration in the review of a proposed essential medicine.
- » Prescriber level
Here the proposed prescriber level should be included. If more than one level is proposed each relevant box should be ticked.

Section 2: Evidence and motivation

- » Estimated benefit
 - Effect measure: this is the clinical outcome that was reported in the clinical trial such as BP, FEV₁, CD₄, VL etc.
 - Risk benefit: this should be reported in the clinical trial and, in most cases, includes the 95% confidence level (95% CI). Absolute risk reduction, also termed risk difference, is the difference between the absolute risk of an event in the intervention group and the absolute risk in the control group.
 - Number Need to Treat (NNT): gives the number of patients who need to be treated for a certain period of time to prevent one event. It is the reciprocal of the absolute risk or can be calculated using the formula below.

Calculations

	Bad outcome	Good outcome	Total patients
Intervention group	<i>a</i>	<i>c</i>	<i>a + c</i>
Control group	<i>b</i>	<i>d</i>	<i>b + d</i>

Measure

Equation

Absolute risk:

$$[b/(b+d)] - [a/(a+c)]$$

Number needed to treat

$$\frac{1}{[b/(b+d)] - [a/(a+c)]}$$

Relative risk

$$[a/(a+c)] \div [b/(b+d)]$$

Odds ratio

$$\frac{[a/(a+c)] \div [c/(a+c)]}{[b/(b+d)] \div [d/(b+d)]} = (a/c) \div (b/d)$$

» Motivating information (**Level of evidence based on the SORT system**)

- The National Essential Drug List Committee has endorsed the adoption of the SORT system for categorising levels of evidence. This system¹ contains only three levels:

Level I	Good quality evidence	Systematic review of RCTs with consistent findings High quality individual RCT
Level II	Limited quality patient orientated evidence	Systematic review of lower quality studies or studies with inconsistent findings Low quality clinical trial Cohort studies Case-control studies
Level III	Other	Consensus guidelines, extrapolations from bench research, usual practice, opinion, disease-oriented evidence (intermediate or physiologic outcomes only), or case series

A: Newer product: for most newer products, level I evidence such as high quality systematic reviews or peer-reviewed high quality randomised controlled trials should be identified and referenced in the space provided.

B: Older products: many of these products were developed prior to the wide use of randomised controlled trials. However, there maybe level I evidence where the product was used as the control arm for a newer product. If no level 1 evidence can be identified, then level II data from poorer quality controlled trials or high quality observational studies should be referenced in the space provided.

» Cost considerations

- Where a published reference supporting the review of cost is available comments should be made regarding its applicability to the South African public sector environment.
- Possible unpublished information that can be included:
 - Cost per daily dose or course of therapy – for long term or chronic therapy such as hypertension the usual daily dose should be calculated (Dose x number of times a day) and converted into the number of dosing units e.g. tablets. This is then used to calculate the cost per day. For medications used in a course of therapy such as antibiotics this is then multiplied by the number of days in the course of therapy.
 - Cost minimisation is used where there is evidence to support equivalence and aims to identify the least costly treatment by identifying all the relevant costs associated with the treatment.
 - Cost-effectiveness analysis is used to compare treatment alternatives that differ in the degree of success in terms of the therapeutic or clinical outcome. By calculating a summary measurement of efficiency (a cost-effectiveness ratio), alternatives with different costs, efficacy rates, and safety rates can be fairly compared along a level playing field.

Where any of these have been performed tick the relevant block and send as an attachment with all the calculations. If possible, the spread sheet should be supplied electronically.

Section 3: Motivator's Details

The receipt of all submission will be acknowledged. In addition, all decisions with supporting arguments will be communicated where appropriate. This section therefore forms a vital link between the motivator and the decision making process.

¹ Ebell MH, Siwek J, Weiss BD, et al. Strength of recommendation taxonomy (SORT): a patient-centered approach to grading evidence in the medical literature. Am Fam Physician. 2004;69:550-6.



DEPARTMENT OF HEALTH
Republic of South Africa

Motivation form for the inclusion of a new medication on the National Essential Medicines List

Section 1: Medication details			
Generic name (or International Non-proprietary Name):			
Proposed indication:			
Prevalence of condition (based on epidemiological data, if any):			
Prescriber level			
Primary Health Care 1	Medical Officer 2	Specialist 3	Designated Specialist 4

Section 2: Evidence and motivation		
2.1 Estimated benefit		
Effect measure		
Risk difference (95% CI)		
NNT		
2.2: Motivating information (Level of evidence based on the SORT system)		
A. Newer product: High quality systematic reviews or peer-reviewed high quality randomised controlled trials (Level I)		
Author	Title	Journal ref
B. Older product with weaker evidence base: Poorer quality controlled trials or high quality observational studies (Level II)		
Author	Title	Journal ref
2.3: Cost-considerations		
Have you worked up the cost?	YES	NO
	Daily cost	Cost-effectiveness analysis
Other relevant cost information if available:		
Author	Title	Journal ref
2.4: Additional motivating comments.		

Section 3: Motivator's Details	
PTC Title:	Date submitted: